

---

<b>Product name:</b>	PCSK9 Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN87597
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant protein of mouse PCSK9
<b>Reactivity:</b>	Mouse
<b>Applications:</b>	WB 1:500-1:2000,ICC/IF 1:100-1:200,FC 1:200-1:500,IP 1:20-1:50
<b>Molecular Weight:</b>	Calculated MW:75 kDa; Observed MW:65 kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	Enables apolipoprotein binding activity; lipoprotein particle binding activity; and low-density lipoprotein particle receptor binding activity. Involved in several processes, including cellular response to insulin stimulus; cellular response to starvation; and regulation of neuron apoptotic process. Acts upstream of or within several processes, including low-density lipoprotein receptor particle metabolic process; regulation of low-density lipoprotein particle receptor catabolic process; and triglyceride metabolic process. Located in COPII-coated ER to Golgi transport vesicle; endoplasmic reticulum; and extracellular space. Is expressed in several structures, including alimentary system; cerebellum; genitourinary system; liver; and telencephalon. Human ortholog(s) of this gene implicated in familial hypercholesterolemia and hypobetalipoproteinemia. Orthologous to human PCSK9 (proprotein convertase subtilisin/kexin type 9). [provided by Alliance of Genome Resources, Apr 2022]