


Product name:	Phospho-AKT1 (S473)
Cat number:	MAB-94621
Conjugate:	Unconjugated
Size:	100 ul
Clone:	587F11
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgGk
Immunogen:	A synthetic Phosphorylated Peptide corresponding to residues target proteinAKT protein
Reactivity:	Hu, Ms, Rt
Applications:	WB 1:1,000 - ICC/IF: 1:200 -1:500 -IHC: 1:100 - 1:200
Molecular Weight:	60 kDa
Purification:	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser473 of human Akt1 protein.
Form:	liquid
Buffer:	Buffer: Supplied in PBS, 50% Glycerol, 0.05% Proclin -For
Storage:	Store: At +4°C for short term, at-20°C for longer term Avoid freezing and thawing cycles

Background:


Background: Akt, also referred to as PKB or Rac, plays a critical role in controlling survival and apoptosis (1-3). This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase (2,3). Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 (4) and by phosphorylation within the carboxy terminus at Ser473. The previously elusive PDK2 responsible for phosphorylation of Akt at Ser473 has been identified as mammalian target of rapamycin (mTOR) in a rapamycin-insensitive complex with rictor and Sin1 (5,6). Akt promotes cell survival by inhibiting apoptosis through phosphorylation and inactivation of several targets, including Bad (7), forkhead transcription factors (8), c-Raf (9), and caspase-9. PTEN phosphatase is a major negative regulator of the PI3 kinase/Akt signaling pathway (10). LY294002 is a specific PI3 kinase inhibitor (11). Another essential Akt function is the regulation of glycogen synthesis through phosphorylation and inactivation of GSK-3 and (12,13). Akt may also play a role in insulin stimulation of glucose transport (12). In addition to its role in survival and glycogen synthesis, Akt is involved in cell cycle regulation by preventing GSK-3-mediated phosphorylation and degradation of cyclin D1 (14) and by negatively regulating the cyclin dependent kinase inhibitors p27 Kip (15) and p21 Waf1/CIP1 (16). Akt also plays a critical role in cell growth by directly phosphorylating mTOR in a rapamycin-insensitive complex containing raptor (17). More importantly, Akt phosphorylates and inactivates tuberin (TSC2), an inhibitor of mTOR within the mTOR-raptor complex (18,19). Specificity/Sensitivity: Phospho-Akt1 (Ser473) (D7F10) Rabbit mAb (Akt1 Specific) recognizes endogenous levels of Akt1 protein only when phosphorylated at Ser473. It does not detect Akt2 protein when phosphorylated at Ser474.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-AKT (Phospho Ser473) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: C6; Lane 2: C6 was treated with IGF-1 (50ng/mL) for 5 minutes-Observed band size: 60kDa



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-AKT (Phospho Ser473) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: NIH-3T3; Lane 2: PC-12-Observed band size: 60kDa



Human lung carcinoma was stained with anti-AKT (Phospho Ser473) rabbit antibody, working conc. 1:100