

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | Phospho-EGFR (Tyr1068) Rabbit Polyclonal Antibody   |
| <b>Cat number:</b>       | ABN03635  |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Polyclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | A synthetic Phosphorylated peptide corresponding to residues target protein   |
| <b>Reactivity:</b>       | Human   |
| <b>Applications:</b>     | WB 1:500-1:1000   |
| <b>Molecular Weight:</b> | Calculated MW: 134 kDa; Observed MW: 175 kDa  |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | EGFR is a receptor tyrosine kinase. Receptor for epidermal growth factor (EGF) and related growth factors including TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. . A single-pass transmembrane tyrosine kinase. Ligand binding to this receptor results in receptor dimerization, autophosphorylation (in trans), activation of various downstream signaling molecules and lysosomal degradation. |