




<b>Product name:</b>	Phospho-MAPKAPK-2 (T334)
<b>Cat number:</b>	ABP-0396
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Poly
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from human MAPKAPK-2 around the phosphorylation site of T334
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:500-1:2000 Immunohistochemistry: 1:100-1:300 ELISA: 1:20000
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using a epitope-specific immunogen.
<b>Form:</b>	liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage:</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Background:</b>	Phospho-MAPKAPK-2 (T334) polyclonal antibody detects endogenous levels of MAPKAPK-2 protein only when phosphorylated at T334.



Western blot analysis of lysates from NIH/3T3 cells, using MAPKAPK2 (Phospho-Thr334) Antibody. The lane on the right is blocked with the phospho peptide



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4\* overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using MAPKAPK2 (Phospho-Thr334) Antibody. The picture on the right is blocked with the phospho peptide.