
Product name:	Phospho-Met (Y1349)
Cat number:	MAB-53309
Conjugate:	Unconjugated
Size:	100 ug
Clone:	13DH2
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Reactivity:	Hu, Ms, Rt
Applications:	WB 1:1000
Molecular Weight:	145 kDa
Purification:	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1349 of human Met.
Form:	liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Storage:	Store at -20°C, and avoid repeat freeze-thaw cycles
Background:	<p>Met, a high affinity tyrosine kinase receptor for hepatocyte growth factor (HGF, also known as scatter factor) is a disulfide-linked heterodimer made of 45 kDa α- and 145 kDa β-subunits (1,2). The α-subunit and the amino-terminal region of the β-subunit form the extracellular domain. The remainder of the β-chain spans the plasma membrane and contains a cytoplasmic region with tyrosine kinase activity. Interaction of Met with HGF results in autophosphorylation at multiple tyrosines, which recruit several downstream signaling components, including Gab1, c-Cbl, and PI3 kinase (3). These fundamental events are important for all of the biological functions involving Met kinase activity. The addition of a phosphate at cytoplasmic Tyr1003 is essential for Met protein ubiquitination and degradation (4). Phosphorylation at Tyr1234/1235 in the Met kinase domain is critical for kinase activation. Phosphorylation at Tyr1349 in the Met cytoplasmic domain provides a direct binding site for Gab1 (5). Altered Met levels and/or tyrosine kinase activities are found in several types of tumors, including renal, colon, and breast. Thus, Met is an attractive cancer therapeutic and diagnostic target (6,7). Phospho-Met (Tyr1349) (130H2) Rabbit mAb detects endogenous levels of Met only when phosphorylated at tyrosine 1349. This antibody may crossreact with other activated protein tyrosine kinases.</p>