

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | PHOX2A Rabbit Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN86388   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | A synthetic peptide of human PHOX2A   |
| <b>Reactivity:</b>       | Human, Mouse  |
| <b>Applications:</b>     | WB 1:1000-1:5000, ICC/IF 1:100-1:200  |
| <b>Molecular Weight:</b> | Calculated MW:30 kDa; Observed MW:36-38 kDa   |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | The protein encoded by this gene contains a paired-like homeodomain most similar to that of the <i>Drosophila aristaless</i> gene product. The encoded protein plays a central role in development of the autonomic nervous system. It regulates the expression of tyrosine hydroxylase and dopamine beta-hydroxylase, two catecholaminergic biosynthetic enzymes essential for the differentiation and maintenance of the noradrenergic neurotransmitter phenotype. The encoded protein has also been shown to regulate transcription of the alpha3 nicotinic acetylcholine receptor gene. Mutations in this gene have been associated with autosomal recessive congenital fibrosis of the extraocular muscles. [provided by RefSeq, Jul 2008] |