

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | Poliovirus Receptor Rabbit Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN03157   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | Recombinant protein of human Poliovirus Receptor  |
| <b>Reactivity:</b>       | Human   |
| <b>Applications:</b>     | WB 1:500-1:1000,IHC 1:50-1:100  |
| <b>Molecular Weight:</b> | Calculated MW: 45 kDa; Observed MW: 60-80 kDa   |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | Mediates NK cell adhesion and triggers NK cell effector functions. Binds two different NK cell receptors: CD96 and CD226. These interactions accumulates at the cell-cell contact site, leading to the formation of a mature immunological synapse between NK cell and target cell. |