

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | Prothrombin Rabbit Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN87330   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | Recombinant protein of mouse Prothrombin  |
| <b>Reactivity:</b>       | Mouse,Rat   |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:200-1:2000  |
| <b>Molecular Weight:</b> | Calculated MW:70 kDa; Observed MW:70 kDa  |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | <p>This gene encodes a vitamin K-dependent glycoprotein coagulation factor that plays an important role in the process of blood coagulation and hemostasis. The encoded protein is an inactive zymogen that undergoes enzymatic cleavage by the coagulation factor Xa to form an active serine protease that converts soluble fibrinogen to insoluble fibrin clot. Most of the mice lacking the encoded protein die at an embryonic stage due to defects in yolk sac vasculature, while the rare neonates succumb to hemorrhage on the first postnatal day.</p> <p>[provided by RefSeq, Apr 2015]</p> |