

---

<b>Product name:</b>	Rab 24 Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN16745
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from Rab 24 . at AA range: 90-170
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>RAB24 is a small GTPase of the Rab subfamily of Ras-related proteins that regulate intracellular protein trafficking (Olkkonen et al., 1993 [PubMed 8126105]).[supplied by OMIM, Aug 2009],function:May be involved in autophagy-related processes.,induction:By extensive retinoic acid treatment, in Ntera-2 teratoma cell line induced to differentiate into post-mitotic neurons (NTN2) (at protein level).,miscellaneous:The unusual Ser-67, instead of a conserved Gln in other family members, is the cause of low GTPase activity. As a result, the predominant nucleotide associated with the protein is GTP.,PTM:Isoprenylation is inefficient compared to other Rab family members.,similarity:Belongs to the small GTPase superfamily. Rab family.,subcellular location:Only about 20-25% is recovered in the particulate fraction.,subunit:Unlike other Rab family members, does not interact with GDP dissociation inhibitors (GDIs), including ARHGDI and ARHGDIIB.,</p>