


Product name:	Rabbit anti mCherry-Tag Monoclonal Antibody
Cat number:	MAB-94804
Conjugate:	Unconjugated
Size:	100 ug
Clone:	125C
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-236 of Discosoma MCherry fluorescent protein.
Reactivity:	Species independent
Applications:	Western Blot: 1:2000 - 1:10000 Immunoprecipitation:0.5µg-4µg antibody for 200µg-600µg extracts of whole cells
Molecular Weight:	55kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.
Storage:	Storage Store at -20°C. Avoid freeze / thaw cycles.
Background:	Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes.Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NTag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.



Western blot analysis of lysates from wild type (WT) and 293F cells transfected with CT55-mCherry-Tag using mCherry-Tag Rabbit mAb at 1:6800 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus. .Exposure time: 20s.



Immunoprecipitation of CT55-mCherry from 300 µg extracts of 293F cells transfected with a CT55 expression vector containing a single C-terminal mCherry-Tag was performed using 3 µg of mCherry-Tag Rabbit mAb. Rabbit IgG isotype control was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using mCherry-Tag Rabbit mAb at a dilution of 1:6000.