

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | Retinoic Acid Receptor gamma Rabbit Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN84272  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthesized peptide derived from human RAR gamma   |
| <b>Reactivity:</b>       | Human,Rat  |
| <b>Applications:</b>     | WB 1:1000-1:2000,ICC 1:50-1:200  |
| <b>Molecular Weight:</b> | 50 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. |