

Product name:	Rhotekin Rabbit Polyclonal Antibody
Cat number:	ABN17137
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from Rhotekin . at AA range: 80-160
Reactivity:	Human,Mouse,Rat
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a scaffold protein that interacts with GTP-bound Rho proteins. Binding of this protein inhibits the GTPase activity of Rho proteins. This protein may interfere with the conversion of active, GTP-bound Rho to the inactive GDP-bound form by RhoGAP. Rho proteins regulate many important cellular processes, including cytokinesis, transcription, smooth muscle contraction, cell growth and transformation. Dysregulation of the Rho signal transduction pathway has been implicated in many forms of cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Mediates Rho signaling to activate NF-kappa-B and may confer increased resistance to apoptosis to cells in gastric tumorigenesis. May play a novel role in the organization of septin structures.,similarity:Contains 1 PH domain.,similarity:Contains 1 REM (Hr1) repeat.,subunit:Interacts via its C-terminal region with the TAX1BP3 PDZ domain. This interaction facilitates Rho-mediated activation of the c-Fos serum response element (SRE). Interacts with SEPT9. Specifically binds to GTP-bound RHOA, RHOB and RHOC and inhibits their GTPase activity.,tissue specificity:Highly expressed in prostate, moderately in kidney, heart, brain, spleen, testis, placenta, small intestine, pancreas, skeletal muscle and peripheral blood leukocytes, and weakly in ovary, colon and thymus. Weakly expressed in all normal cell lines tested. Overexpressed in various cancer cell lines.,