
Product name:	ST6GAL1 Rabbit Polyclonal Antibody
Cat number:	ABN18325
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthetic peptide from human protein at AA range: 63-135
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	<p>This gene encodes a member of glycosyltransferase family 29. The encoded protein is a type II membrane protein that catalyzes the transfer of sialic acid from CMP-sialic acid to galactose-containing substrates. The protein, which is normally found in the Golgi but can be proteolytically processed to a soluble form, is involved in the generation of the cell-surface carbohydrate determinants and differentiation antigens HB-6, CD75, and CD76. This gene has been incorrectly referred to as CD75. Three transcript variants encoding two different isoforms have been described. [provided by RefSeq, Aug 2009],catalytic activity:CMP-N-acetylneuraminate + beta-D-galactosyl-1,4-N-acetyl-beta-D-glucosamine = CMP + alpha-N-acetylneuraminy-2,6-beta-D-galactosyl-1,4-N-acetyl-beta-D-glucosamine.,function:Transfers sialic acid from the donor of substrate CMP-sialic acid to galactose containing acceptor substrates.,online information:GlycoGene database,online information:ST6Gal I,pathway:Protein modification; protein glycosylation.,PTM:The HB-6, CDW75, and CD76 differentiation antigens are cell-surface carbohydrate determinants generated by this enzyme.,PTM:The soluble form derives from the membrane form by proteolytic processing.,similarity:Belongs to the glycosyltransferase 29 family.,subcellular location:Membrane-bound form in trans cisternae of Golgi. Secreted into the body fluid.,</p>