

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | TCP 1 alpha Rabbit Monoclonal antibody   |
| <b>Cat number:</b>       | MABN03224  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthetic peptide of human TCP1 alpha/CCTA   |
| <b>Reactivity:</b>       | Human,Hamster  |
| <b>Applications:</b>     | WB 1:500-1:1000,IHC 1:50-1:100,ICC/IF 1:50-1:200,IP 1:20-1:50  |
| <b>Molecular Weight:</b> | Calculated MW: 60 kDa; Observed MW: 60 kDa   |
| <b>Purification:</b>     | Affinity Purified  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Molecular chaperone; assists the folding of proteins upon ATP hydrolysis. As part of the BBS/CCT complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia. Known to play a role, in vitro, in the folding of actin and tubulin. |