

<b>Product name:</b>	TGF beta Receptor II (17K1) Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN18849
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant protein of human TGF beta Receptor II
<b>Reactivity:</b>	Human
<b>Applications:</b>	ICC/IF 1:500-1:1000,FC 1:20-1:50
<b>Molecular Weight:</b>	65kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**Background:**

TGFβs mediate their activity by high affinity binding to the type II receptor (TGFβ RII) transmembrane protein with a cytoplasmic serine-threonine kinase domain. For signaling growth inhibition and early gene responses the type II receptor requires both its kinase activity and association with a TGFβ-binding protein, designated the type I receptor. Two independent groups have recently described the cloning and sequence analysis of genes encoding TGFβ type I receptor proteins designated ALK-5 (TβR-1) and TSR-1, respectively. Transmembrane serine/threonine kinase forming with the TGF- beta type I serine/threonine kinase receptor, TGFBR1, the non- promiscuous receptor for the TGF-beta cytokines TGFβ1, TGFβ2 and TGFβ3. Transduces the TGFβ1, TGFβ2 and TGFβ3 signal from the cell surface to the cytoplasm and is thus regulating a plethora of physiological and pathological processes including cell cycle arrest in epithelial and hematopoietic cells, control of mesenchymal cell proliferation and differentiation, wound healing, extracellular matrix production, immunosuppression and carcinogenesis. The formation of the receptor complex composed of 2 TGFBR1 and 2 TGFBR2 molecules symmetrically bound to the cytokine dimer results in the phosphorylation and the activation of TGFBR1 by the constitutively active TGFBR2. Activated TGFBR1 phosphorylates SMAD2 which dissociates from the receptor and interacts with SMAD4. The SMAD2-SMAD4 complex is subsequently translocated to the nucleus where it modulates the transcription of the TGF-beta-regulated genes. This constitutes the canonical SMAD-dependent TGF-beta signaling cascade. Also involved in non-canonical, SMAD-independent TGF-beta signaling pathways.