
Product name:	TPA Tissue Plasminogen Activator Rabbit Monoclonal Antibody
Cat number:	MABN86548
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human TPA Tissue Plasminogen Activator
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:1000-1:5000,IHC 1:100-1:1000
Molecular Weight:	Calculated MW:63 kDa; Observed MW:63 kDa
Purification:	Affinity Purification
Form:	Liquid
Buffer:	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	This gene encodes tissue-type plasminogen activator, a secreted serine protease that converts the proenzyme plasminogen to plasmin, a fibrinolytic enzyme. The encoded preproprotein is proteolytically processed by plasmin or trypsin to generate heavy and light chains. These chains associate via disulfide linkages to form the heterodimeric enzyme. This enzyme plays a role in cell migration and tissue remodeling. Increased enzymatic activity causes hyperfibrinolysis, which manifests as excessive bleeding, while decreased activity leads to hypofibrinolysis, which can result in thrombosis or embolism. Alternative splicing of this gene results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016]