

---

<b>Product name:</b>	TWIK-1 Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN19444
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human KCNK1. AA range:287-336
<b>Reactivity:</b>	Human,Rat
<b>Applications:</b>	WB 1:500-1:2000,ELISA 1:20000-1:40000
<b>Molecular Weight:</b>	38kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	This gene encodes one of the members of the superfamily of potassium channel proteins containing two pore-forming P domains. The product of this gene has not been shown to be a functional channel, however, it may require other non-pore-forming proteins for activity. [provided by RefSeq, Jul 2008],function:Weakly inward rectifying potassium channel.,miscellaneous:Inhibited by barium, quinine, quinidine and internal acidification. Activated by protein kinase C.,similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family.,subunit:Homodimer .,tissue specificity:Widely expressed with high levels in heart and brain and lower levels in placenta, lung, liver and kidney.,