

Product name:	USP16 Rabbit Polyclonal Antibody
Cat number:	ABN19666
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from the Internal region of human USP16.
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight:	93kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a deubiquitinating enzyme that is phosphorylated at the onset of mitosis and then dephosphorylated at the metaphase/anaphase transition. It can deubiquitinate H2A, one of two major ubiquitinated proteins of chromatin, in vitro and a mutant form of the protein was shown to block cell division. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008], catalytic activity: Ubiquitin C-terminal thioester + H₂O = ubiquitin + a thiol., disease: A chromosomal aberration involving USP16 is a cause of Chronic myelomonocytic leukemia. Inversion inv(21) (q21;q22) with RUNX1/AML1., domain: The UBP-type zinc finger binds 3 zinc ions that form a pair of cross-braced ring fingers encapsulated within a third zinc finger in the primary structure. It recognizes the C-terminal tail of free ubiquitin., function: Specifically deubiquitinates histone H2A, a specific tag for epigenetic transcriptional repression, thereby acting as a coactivator. Deubiquitination of histone H2A is a prerequisite for subsequent phosphorylation at 'Ser-10' of histone H3, and is required for chromosome segregation when cells enter into mitosis. Regulates Hox gene expression via histone H2A deubiquitination. Prefers nucleosomal substrates. Does not deubiquitinate histone H2B., PTM: Phosphorylated at the onset of mitosis and dephosphorylated during the metaphase/anaphase transition. The phosphorylated form of the protein is also enzymatically active., similarity: Belongs to the peptidase C19 family. USP16 subfamily., similarity: Contains 1 UBP-type zinc finger., subunit: Homotetramer., tissue specificity: Present in all the tissues examined including fetal brain, lung, liver, kidney, and adult heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.,