
| | |
|--------------------------|--|
| Product name: | Vinculin Rabbit Monoclonal Antibody |
| Cat number: | MABN02763 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | Recombinant protein of human Vinculin |
| Reactivity: | Human, Mouse, Rat |
| Applications: | WB 1:500-1:1000, ICC/IF 1:50-1:200, IP 1:20-1:50 |
| Molecular Weight: | Calculated MW: 124 kDa; Observed MW: 124 kDa |
| Purification: | Affinity Purification |
| Form: | Liquid |
| Buffer: | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | Vinculin is a cytoskeletal protein that plays an important role in the regulation of focal adhesions and embryonic development. Three structural vinculin domains include an amino-terminal head, a short, flexible proline-rich region and a carboxy-terminal tail. In the inactive state, the head and tail domains of vinculin interact to form a closed conformation. The open and active form of vinculin translocates to focal adhesions where it is thought to be involved in anchoring F-actin to the membrane and regulation of cell migration. |