

---

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | WDHD1 Rabbit Polyclonal Antibody   |
| <b>Cat number:</b>       | ABN19874   |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Polyclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human WDHD1. AA range:721-770  |
| <b>Reactivity:</b>       | Human,Rat,Mouse  |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000   |
| <b>Molecular Weight:</b> | 126kDa   |
| <b>Purification:</b>     | Affinity purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | <p>The protein encoded by this gene contains multiple N-terminal WD40 domains and a C-terminal high mobility group (HMG) box. WD40 domains are found in a variety of eukaryotic proteins and may function as adaptor/regulatory modules in signal transduction, pre-mRNA processing and cytoskeleton assembly. HMG boxes are found in many eukaryotic proteins involved in chromatin assembly, transcription and replication. Alternative splicing results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Binds DNA with high affinity. May also be involved in protein-protein interactions.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 HMG box DNA-binding domain.,similarity:Contains 7 WD repeats.,</p> |