

---

<b>Product name:</b>	WRN Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN86692
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthetic peptide of human WRN
<b>Reactivity:</b>	Human
<b>Applications:</b>	WB 1:1000-1:5000
<b>Molecular Weight:</b>	Calculated MW:163 kDa; Observed MW:200 kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	This gene encodes a member of the RecQ subfamily of DNA helicase proteins. The encoded nuclear protein is important in the maintenance of genome stability and plays a role in DNA repair, replication, transcription and telomere maintenance. This protein contains a N-terminal 3' to 5' exonuclease domain, an ATP-dependent helicase domain and RQC (RecQ helicase conserved region) domain in its central region, and a C-terminal HRDC (helicase RNase D C-terminal) domain and nuclear localization signal. Defects in this gene are the cause of Werner syndrome, an autosomal recessive disorder characterized by accelerated aging and an elevated risk for certain cancers. [provided by RefSeq, Aug 2017]