

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | Xanthine Oxidase Rabbit Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN02776  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthetic peptide of human Xanthine Oxidase  |
| <b>Reactivity:</b>       | Human  |
| <b>Applications:</b>     | WB 1:500-1:1000  |
| <b>Molecular Weight:</b> | Calculated MW: 146 kDa; Observed MW: 146 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Key enzyme in purine degradation. Catalyzes the oxidation of hypoxanthine to xanthine. Catalyzes the oxidation of xanthine to uric acid. Contributes to the generation of reactive oxygen species. Has also low oxidase activity towards aldehydes (in vitro). |