

---

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | ZNF75 Rabbit Polyclonal Antibody   |
| <b>Cat number:</b>       | ABN20284   |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Polyclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human ZNF75. AA range:296-345  |
| <b>Reactivity:</b>       | Human,Rat,Mouse  |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000   |
| <b>Molecular Weight:</b> | 40kDa  |
| <b>Purification:</b>     | Affinity purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | <p>zinc finger protein 75D(ZNF75D) Homo sapiens This gene encodes a protein that likely functions as a transcription factor. The protein, which belongs to the ZNF75 family, includes an N-terminal SCAN domain, a KRAB box, and five C2H2-type zinc finger motifs. Another functional gene belonging to this family is located on chromosome 16, while pseudogenes have been identified on chromosomes 11 and 12. Alternative splicing results in multiple transcripts variants. [provided by RefSeq, Jun 2010],function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 KRAB domain.,similarity:Contains 1 SCAN box domain.,similarity:Contains 5 C2H2-type zinc fingers.,</p> |